

Clotrizen

Clotrimazole USP 1%

Composition

Clotrizen Cream: Each gram cream contains Clotrimazole USP 10 mg.

Pharmacology

Clotrimazole is a broad spectrum antifungal. It also exhibits activity against Trichomonas, Staphylococci, Streptococci and Bacteroides

Indication

1. All dermatomycoses due to dermatophytes (e.g. Trichophyton species).
2. All dermatomycoses due to yeasts (Candida species).
3. Dermatomycoses due to moulds and other fungi.
4. Skin diseases showing superinfections with these fungi.

Some examples of skin infections included in 1-4 points are: interdigital mycoses (e.g. athlete's foot), paronychias (associated with nail mycoses), mycoses in skin folds, Candida vulvitis, Candida balanitis, Pityriasis versicolor, erythrasma.

Dosage and Administration

Clotrizen Cream should be thinly applied 2-3 times daily to the infected sites and rubbed in. On account of the excellent efficacy, a small amount of cream is usually sufficient for an area about the size of the palm. For the treatment to become a complete success, reliable and sufficiently long time application of Clotrizen Cream is important. The duration of treatment varies; it depends among other factors on the extent and localisation of the disease.

Recommended Duration Of Treatment

Dermatomycoses : 3-4 week

Candida vulvitis and Candida balanitis: 1-2 weeks

Erythrasma and Pityriasis versicolor :3 weeks (approximately).

In fungal infection of the feet, to prevent relapse, treatment should be continued for about 2 weeks

beyond the disappearance of all signs of disease. After washing, the feet should be thoroughly dried (particularly spaces between the toes). Clotrizen Cream is odorless, can be washed off and does not stain clothing.

Contraindication & Precaution

Hypersensitivity to Clotrimazole

Side effects

When applied topically, Clotrimazole is well tolerated. With external application, systemic effects are not observed. Local irritation or burning sensation may occur in a very few cases but these symptoms are not considered harmful.

Use in Pregnancy & Lactation

It is recommended that Clotrimazole should be used in pregnancy only when considered necessary by the physician.

Drug interaction

No information is available

Storage

Store below 30°C in a dry place, away from light. Keep out of the reach of children.

Packaging

Clotrizen Cream: Each pack has a laminated tube containing 10 gm cream.

Manufactured By

BEACON[®]
Pharmaceuticals PLC

Bhaluka, Mymensingh, Bangladesh