

# Resva

Rosuvastatin

## COMPOSITION

**Resva 5 Tablet:** Each film coated tablet contains Rosuvastatin Calcium BP equivalent to 5 mg Rosuvastatin.

**Resva 10 Tablet:** Each film coated tablet contains Rosuvastatin Calcium BP equivalent to 10 mg Rosuvastatin.

**Resva 20 Tablet:** Each film coated tablet contains Rosuvastatin Calcium BP equivalent to 20 mg Rosuvastatin.

## PHARMACOLOGY

**Resva** (Rosuvastatin) is a selective and competitive inhibitor of HMG-CoA reductase, the rate-limiting enzyme that converts 3-hydroxy-3-methylglutaryl coenzyme A to mevalonate, a precursor of cholesterol. Rosuvastatin produces its lipid-modifying effects in two ways. First, it increases the number of hepatic LDL receptors on the cell surface to enhance uptake and catabolism of LDL. Second, Rosuvastatin inhibits hepatic synthesis of VLDL, which reduces the total number of VLDL and LDL particles.

## PHARMACOKINETICS

### Absorption

Peak plasma concentrations of Rosuvastatin were reached 3 to 5 hours following oral dosing. Both peak concentration (C<sub>max</sub>) and area under the plasma concentration-time curve (AUC) increased in approximate proportion to Rosuvastatin dose.

### Distribution

Mean volume of distribution at steady-state of Rosuvastatin is approximately 134 liters. Rosuvastatin is 88% bound to plasma proteins, mostly albumin.

### Metabolism

Rosuvastatin is not extensively metabolized. The major metabolite of Rosuvastatin is N-desmethyl rosuvastatin which has approximately one-sixth to one-half the HMG-CoA reductase inhibitory activity of Rosuvastatin. Overall, greater than 90% of active plasma HMG-CoA reductase inhibitory activity is accounted for by Rosuvastatin.

### Elimination

Following oral administration, Rosuvastatin and its metabolites are primarily excreted in the feces (90%). The elimination half-life (t<sub>1/2</sub>) of Rosuvastatin is approximately 19 hours.

## INDICATION AND USE

- Heterozygous Hypercholesterolemia (Familial and Nonfamilial)
- Homozygous Hypercholesterolemia (Familial)
- Mixed Dyslipidemia (Fredrickson Type IIa and IIb)

## DOSAGE AND ADMINISTRATION

*Heterozygous Hypercholesterolemia (Familial and Nonfamilial) and Mixed Dyslipidemia (Fredrickson Type IIa and IIb)*

The usual recommended starting dose of Rosuvastatin is 10 mg once daily. Initiation of therapy with 5 mg once daily may be considered for patients requiring less aggressive LDL-C reductions or who have predisposing factors for myopathy. For patients with marked hypercholesterolemia (LDL-C > 190 mg/dL) and aggressive lipid targets, a 20 mg starting dose may be considered. The 40 mg dose of Rosuvastatin should be reserved for those patients who have not achieved goal LDL-C at 20 mg. After initiation and/or upon titration of Rosuvastatin, lipid levels should be analyzed within 2 to 4 weeks and dosage adjusted accordingly.

### *Homozygous Hypercholesterolemia (Familial)*

The recommended starting dose of Rosuvastatin is 20 mg once daily in patients with homozygous FH. The maximum recommended daily dose is 40 mg. Rosuvastatin should be used in these patients as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable. Patients with renal insufficiency No modification of dosage is necessary for patients with mild to moderate renal insufficiency. For patients with severe renal impairment (CL<sub>cr</sub> <30 mL/min/1.73 m<sup>2</sup>) not on hemodialysis, dosing of Rosuvastatin should be started at 5 mg once daily and not to exceed 10 mg once daily.

### *Dosage in Asian Patients*

35 Initiation of Rosuvastatin therapy with 5 mg once daily should be considered for Asian patients.

*Use with Cyclosporine, Lopinavir/Ritonavir or Atazanavir/Ritonavir*

In patients taking Cyclosporine, the dose of Rosuvastatin should be limited to 5 mg once daily. In patients taking Lopinavir and Ritonavir or Atazanavir and Ritonavir the dose of Rosuvastatin should be limited to 10 mg once daily.

## CONTRAINDICATION

Rosuvastatin is contraindicated in patients with a known hypersensitivity to any component of this product. Rosuvastatin is contraindicated in patients with active liver disease or with unexplained persistent elevations of serum transaminases.

## SIDE EFFECT

Rosuvastatin is generally well tolerated. The most frequent adverse events thought to be related to Rosuvastatin were myalgia, constipation, asthenia, abdominal pain, and nausea.

## WARNING

In case of myopathy, rhabdomyolysis, acute renal failure; Please consult with physician.

## DRUG INTERACTION

**Erythromycin:** Co-administration of Erythromycin with Rosuvastatin decreased AUC and C<sub>max</sub> of Rosuvastatin by 20% and 31%, respectively. **Itraconazole:** Itraconazole resulted in a 39% and 28% increase in AUC of Rosuvastatin after 10 mg and 80 mg dosing, respectively. **Fluconazole:** Co-administration of Fluconazole with Rosuvastatin resulted in a 14% increase in AUC of Rosuvastatin. **Warfarin:** Co-administration of Warfarin (25 mg) with Rosuvastatin (40 mg) did not change Warfarin plasma concentrations but increased the International Normalized Ratio (INR). **Gemfibrozil:** Co-administration of Gemfibrozil (600 mg twice daily for 7 days) with Rosuvastatin (80 mg) resulted in a 90% and 120% increase for AUC and C<sub>max</sub> of Rosuvastatin, respectively. **Antacid:** Co-administration of an antacid (aluminium and magnesium hydroxide combination) with Rosuvastatin (40 mg) resulted in a decrease in plasma concentrations of Rosuvastatin by 54%. **Oral contraceptives:** Co-administration of oral contraceptives (Ethinyl Estradiol and Norgestrel) with Rosuvastatin resulted in an increase in plasma concentrations of Ethinyl Estradiol and Norgestrel by 26% and 34%, respectively.

## USE IN PREGNANCY AND LACTATION

Pregnancy Category: X

## STORAGE CONDITION

Store in a cool (below 30°C) and dry place, away from light. Store in the original package in order to protect from moisture. Keep out of the reach of children.

## Commercial pack

**Resva 5 Tablet:** Each commercial box contains 28 film coated tablets in Alu-Alu blister pack.

**Resva 10 Tablet:** Each commercial box contains 28 film coated tablets in Alu-Alu blister pack.

**Resva 20 Tablet:** Each commercial box contains 14 film coated tablets in Alu-Alu blister pack.

Manufactured By  
 **BEACON**<sup>®</sup>  
Pharmaceuticals Limited  
Bhaluka, Mymensingh, Bangladesh