

# Art work of Tivis Insert



## Composition

**Tivis Tablet:** Each film-coated tablet contains Tiemonium Methylsulphate INN 50 mg.

**Tivis injection:** Each 2 ml ampoule contains Tiemonium Methylsulphate INN 5 mg

## Pharmacological action

Tiemonium Methylsulphate is a synthetic antispasmodic agent. Tiemonium strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract

## Indications

Tiemonium Methylsulphate is an antispasmodic drug that reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal, biliary, urological and gynecological disease such as gastroenteritis, diarrhoea, dysentery, biliary colic, cholecystitis, colonopathies, mild cystitis, & spasmodic dysmenorrhoea. Tiemonium Methylsulphate IM/IV Injection is indicated for acute pain.

## Dosage and Administration

### Oral

**Tablet:** 2 to 6 tablets (100 - 300 mg) daily in divided doses as required.

### Parenteral

The usual dose is 5 mg / 2 ml (1 ampoule) three times daily by slow IV or IM injection.

## Side Effects

As with any active substance this drug may cause certain undesirable or unpleasant effects. In some people dry mouth with difficulty in swallowing and thirst, dilatation of pupils with loss of accommodation and sensitivity to light, increased intraocular pressure, flushing, dry skin, bradycardia followed by tachycardia, palpitations and arrhythmias, difficulty in micturition and constipation may occur.

## Contraindications

Tiemonium Methylsulphate should not be used in glaucoma (acute pain of eyeball with vision disturbances) and in difficulty to urinate (disorders of prostate or bladder).

## Overdose

There is not available data regarding the overdose of Tiemonium Methylsulphate.

## Drug interactions

Tiemonium Methylsulphate tablet should not be used with other drugs without prior consult of a registered physician to avoid possible drug interaction.

## Use in Pregnancy & lactation

The use of Tiemonium Methylsulphate is not recommended during pregnancy and lactation, unless the potential benefit outweighs the possible risks.

## Pharmaceutical Information

### Storage condition

Store in a cool and dry place, away from light. Keep out of the reach of children.

### Presentation & Packaging

**Tivis Tablet:** Each commercial box contains 50 tablets in blister pack.

**Tivis injection:** Each commercial box contains 5 ampoules in blister pack.

Manufactured By

**BEACON**<sup>®</sup>  
Pharmaceuticals PLC

Bhaluka, Mymensingh, Bangladesh

LF08101

|       |       |        |        |
|-------|-------|--------|--------|
| Width | 86 mm | Length | 244 mm |
|-------|-------|--------|--------|

# Tivis

Tiemonium Methylsulphate

## Composition

**Tivis Tablet:** Each film-coated tablet contains Tiemonium Methylsulphate INN 50 mg.

**Tivis injection:** Each 2 ml ampoule contains Tiemonium Methylsulphate INN 5 mg

## Pharmacological action

Tiemonium Methylsulphate is a synthetic antispasmodic agent. Tiemonium strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract

## Indications

Tiemonium Methylsulphate is an antispasmodic drug that reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal, biliary, urological and gynecological disease such as gastroenteritis, diarrhoea, dysentery, biliary colic, cholecystitis, colonopathies, mild cystitis, & spasmodic dysmenorrhoea. Tiemonium Methylsulphate IM/IV Injection is indicated for acute pain.

## Dosage and Administration

### Oral

*Tablet:* 2 to 6 tablets (100 - 300 mg) daily in divided doses as required.

### Parenteral

The usual dose is 5 mg / 2 ml (1 ampoule) three times daily by slow IV or IM injection.

## Side Effects

As with any active substance this drug may cause certain undesirable or unpleasant effects. In some people dry mouth with difficulty in swallowing and thirst, dilatation of pupils with loss of accommodation and sensitivity to light, increased intraocular pressure, flushing, dry skin, bradycardia followed by tachycardia, palpitations and arrhythmias, difficulty in micturition and constipation may occur.

## Contraindications

Tiemonium Methylsulphate should not be used in glaucoma (acute pain of eyeball with vision disturbances) and in difficulty to urinate (disorders of prostate or bladder).

## Overdose

There is not available data regarding the overdose of Tiemonium Methylsulphate.

## Drug interactions

Tiemonium Methylsulphate tablet should not be used with other drugs without prior consult of a registered physician to avoid possible drug interaction.

## Use in Pregnancy & lactation

The use of Tiemonium Methylsulphate is not recommended during pregnancy and lactation, unless the potential benefit outweighs the possible risks.

## Pharmaceutical Information

### Storage condition

Store in a cool and dry place, away from light. Keep out of the reach of children.

## Presentation & Packaging

**Tivis Tablet:** Each commercial box contains 50 tablets in blister pack.

**Tivis injection:** Each commercial box contains 5 ampoules in blister pack.

Manufactured By

**BEACON**®

Pharmaceuticals PLC

Bhaluka, Mymensingh, Bangladesh